

SEMIPRIME ANTI FLEXIBLE RINGS

C. JAYA SUBBA REDDY¹ & K. CHENNAKESAVULU²

¹Department of Mathematics, S.V. University, Tirupati, Andhra Pradesh, India

²Department of Mathematics, Intel Engineering College, Anantapur, Andhra Pradesh, India

ABSTRACT

In this paper we prove that the commutator is in the middle nucleus of an anti flexible ring and an anti flexible ring without non-zero nilpotent elements is flexible. Using these properties to show that a semi prime anti flexible ring is weakly standard ring. At the end of this paper we give an example of an anti flexible ring which is not a weakly standard ring.

KEYWORDS: Commutator, Nucleus, Anti Flexible Ring, Flexible Ring, Semi Prime, Weakly Standard Ring

INTRODUCTION

Anti flexible rings were introduced by Koiser [3] and a subclass of anti flexible rings was studied by Klien fled [2]. Simple anti flexible rings are characterized by Anderson and Outcalt [1]. In this paper, we discuss some properties of an anti flexible ring R . In general an anti flexible ring is not a weakly standard ring. We prove that the commutator is in the middle nucleus and an anti flexible ring without non-zero nilpotent elements is flexible. We use these properties to show that R is a weakly standard ring. Also we present an example of an anti flexible ring which is not a weakly standard ring.

PRELIMINARIES

A weakly standard ring R , where the associator $(x, y, z) = (xy)z - x(yz)$ and the commutator $(x, y) = xy - yx$. The nucleus N of R is defined as the set of all elements n in R such that $(n, R, R) = (R, n, R) = (R, R, n) = 0$. We define a ring R is semi prime if for any ideal A of R , $A^2 = 0$ implies $A = 0$. We know that the ring R is said to be anti flexible if

$$A(x, y, z) = (x, y, z) - (z, y, x) = 0 \quad (1)$$

Throughout this paper we assume that R is an anti flexible ring of

$$\text{Characteristic} \neq 2 \text{ and that } (x, x, x) = 0 \quad (2)$$

Is an identity in R . Using the identity (1) in the linearization of (2), we obtain the identity

$$B(x, y, z) = (x, y, z) + (y, z, x) + (z, x, y) = 0 \quad (3)$$

We shall also require the Teichmuller identity (which holds in any ring):

$$C(w, x, y, z) = (wx, y, z) - (w, xy, z) + (w, x, yz) - w(x, y, z) - (w, x, y)z = 0 \quad (4)$$

In any ring $(xy, z) = x(y, z) + (x, z)y + (x, y, z) + (z, x, y) - (x, z, y)$, from which we subtract

$$A(x, z, y) + B(x, y, z) = 0 \text{ to obtain}$$

$$D(x, y, z) = (xy, z) - x(y, z) - (x, z)y + 2(x, z, y) = 0 \quad (5)$$

Expanding $0 = C(w, x, y, z) - C(z, y, x, w)$, and using

$$0 = A(z, y, xw) = A(z, yx, w) = A(zy, x, w) = ZA(y, x, w) = A(z, y, x) w,$$

We get

$$0 = E(w, x, y, z) = ((w, x), y, z) - (w, (x, y), z) + (w, x, (y, z)) - (w, (x, y, z)) - ((w, x, y), z). \quad (6)$$

Then we expand

$$0 = E(w, x, y, z) + E(x, y, z, w) + E(y, z, w, x) + E(z, w, x, y)$$

$$-B((w, x), y, z) - B((x, y), z, w) - B((y, z), w, x) - B((z, w), x, y) \text{ to get}$$

$$0 = F(w, x, y, z) = (w, (x, y), z) + (x, (y, z), w) + (y, (z, w), x) + (z, (w, x), y) \quad (7)$$

$$\text{Now we are able to derive the important identity } (w, (x, y), z) = 0 \quad (8)$$

Using these identities first we prove the following lemmas.

MAIN RESULTS

Lemma 1: *In an anti flexible ring R , commutators are in the middle nucleus.*

Proof: By expanding

$$0 = E(x, x, y, x) + E(y, x, x, x) - B(x, x, (y, x)) + (B(x, x, y), x), \text{ we get}$$

$$0 = (x, (x, y, x)), \text{ hence from } 0 = (x, B(x, y, x)) \text{ and } 0 = (x, A(x, x, y)), \text{ we have}$$

$$0 = (x, (x, y, x)) = (x, (x, x, y)) = (x, (y, x, x)) \quad (9)$$

Then it follows from (9) and $0 = E(y, x, x, x)$ that $((y, x), x, x) = 0$. Hence $(x, x, (y, x)) = 0$ by (1) and then from $0 = B(x, x, (y, x))$, we have

$$0 = (x, (y, x), x) \quad (10)$$

Substituting $x + z$ for x in (10) and subtracting

$$0 = A(z, (y, x), x) + A(z, (y, z), x), \text{ we obtain}$$

$$2(x, (y, x), z) + 2(x, (y, z), z) + (x, (y, z), x) + (z, (y, x), z) = 0 \quad (11)$$

Substituting $-z$ for z in (11) and then adding to (11) yields

$$2(x, (y, z), z) + (z, (y, x), z) = 0 \quad (12)$$

Next we linearize (12) and add $A(w, (y, x), z) = 0$ to get

$$G(w, x, y, z) = (x, (y, z), w) + (x, (y, w), z) + (w, (y, x), z) = 0.$$

Computing

$$0 = F(w, x, y, z) + G(w, x, y, z) + G(x, y, w, z) - A(z, (w, x), y), \text{ we get}$$

$$H(w, x, y, z) = (x, (y, z), w) + (y, (w, x), z) = 0.$$

Now, identity (8) follows from the expansion of

$$0 = H(w, x, y, z) + H(x, w, y, z) - A(w, (y, z), x) 0$$

Lemma 2: In an anti flexible ring, $(x, y, x)^2 = 0$.

Proof: By forming $0 = C(w, x, y, z) - C(x, y, z, w) + C(y, z, w, x) - C(z, w, x, y)$, and using (3),

We obtain

$$I(w, x, y, z) = (w, (x, y, z)) - (x, (y, z, w)) + (y, (z, w, x)) - (z, (w, x, y)) = 0 \quad (13)$$

If in (9) we replace x by both $x + z$ and $-x + z$, adds and divides by 2, we obtain

$$(z, (x, y, x)) + (x, (z, y, x)) + (x, (x, y, z)) = 0,$$

$$(z, (x, x, y)) + (x, (z, x, y)) + (x, (x, z, y)) = 0 \quad (14)$$

$$\text{And } (z, (y, x, x)) + (x, (y, z, x)) + (x, (y, x, z)) = 0.$$

Because of (14) we have

$$\begin{aligned} 0 &= I(x, y, x, z) \\ &= (x, (y, x, z)) - (y, (x, z, x)) + (x, (z, x, y)) - (z, (x, y, x)) \\ &= (x, (y, x, z)) - (y, (x, z, x)) - (z, (x, x, y)) - (x, (x, z, y)) + (x, (z, y, x)) + (x, (x, y, z)) \\ &= 2(x, (x, y, z)) + (x, (y, x, z)) + (x, (x, z, y)) \\ &= (x, (x, y, z)) + (y, z, x) + (z, x, y) + (x, (x, y, z)) = 0. \end{aligned}$$

$$\text{Consequently } (x, (x, y, z)) = 0 \quad (15)$$

By linearizing (15) we obtain

$$J(w, x, y, z) = (w, (x, y, z)) + (x, (w, y, z)) = 0 \quad (16)$$

We define u by $u = (x, y, x)$. Then $0 = J(xy, x, y, x)$ implies $-(xy, u) = (x, (xy, y, x))$.

If we take $z = xy$ in (15), $(x, (x, y, xy)) = 0$ implies $(x, (xy, y, x)) = 0$, using (1). Therefore $(xy, u) = 0$. Because of (9) we have $(u, x) = 0$ and $((y, x, x), y) = 0$. But then $(B(y, x, x) + A(x, y, x) -$

$$2(y, x, x), y) = 0 \text{ implies } (u, y) = 0. \text{ From (5), by taking } z = u, \text{ we get } (x, u, y) = 0.$$

$$\text{Using (3) we obtain } (x, y, u) = -(u, x, y) = (y, u, x) = 0.$$

$$\text{Thus } (u, x, y) = 0 \quad (17)$$

If we put $x = u, y = x, z = x$ in (5) we get $(ux, x) = 0$.

$$\text{By linearizing this, } (ux, y) + (uy, x) = 0.$$

$$\text{Then (5) implies that } ((ux) y, x) = ux(y, x) + 2(ux, y, x).$$

$$\text{By interchanging } x \text{ and } y \text{ in this equation, we have } ((ux) x, y) = ux(x, y) + 2(ux, x, y).$$

By adding these two equations, we obtain

$$(ux, y, x) + (ux, x, y) = 0 \quad (18)$$

From (3), $(ux, y, x) + (y, x, ux) + (x, ux, y) = 0$.

Then $-(ux, x, y) + (y, x, ux) + (x, ux, y) = 0$, using (18).

This becomes $-(y, x, ux) + (y, x, ux) + (x, ux, y) = 0$, using (1).

So $(x, ux, y) = 0$. Also (18) implies that $(ux, y, x) = -(y, x, ux) = (x, ux, y) = 0$. (19)

By using (19) and (17) in (4), we get that $u(x, y, x) = 0$ and hence $(x, y, x)^2 = 0$.

Theorem 1: If R is an anti flexible ring without non-zero nilpotent elements, then R is a weakly standard ring.

Proof: Since R is without nonzero nilpotent elements, lemma (2) implies that

$$(x, y, x) = 0 \quad (20)$$

So R is flexible. Since R is flexible, we have $(x, y, z) = -(z, y, x)$.

By taking $z = v$ as a commutator, we obtain $(x, y, v) = -(v, y, x)$. From (1), it follows that

$$2(x, y, v) = 0 \text{ and hence } (x, y, v) = 0 \quad (21)$$

Since R is of characteristic $\neq 2$. Now from (20), lemma (1) and (21), we get that R is a

Weakly standard ring

Now we give an example of an anti flexible ring which is not a weakly standard ring.

Example 1: Suppose that the ring R is defined by the following multiplication table together with all finite sums of e, a, b, c, d, h such that $x + x = 2x \neq 0$.

	e	a	b	c	d	h
e	e	b	a	0	0	0
a	h	c	0	0	0	0
b	0	0	0	0	0	0
c	0	0	0	0	0	0
d	0	0	0	0	0	0
h	h+b	b	0	0	0	0

We observe that $(a, a, a) = a^2 a - a a^2 = ca - ac = 0$.

Therefore, R satisfies $(x, x, x) = 0$.

This is enough to check the identity $(x, y, z) = (z, y, x)$.

Now $(e, e, a) = ea - e(ea) = b$ and

$$(a, e, e) = (ae)e - ae = he - h = h + b - h = b.$$

Thus $(e, e, a) = (a, e, e)$.

Next we see that the commutator is not in the left nucleus.

$$((e, b), e, e) = (eb, e, e) - (be, e, e)$$

$$= (a, e, e) = b \neq 0.$$

Hence R is not a weakly standard ring.

CONCLUSIONS

In this paper we proved that the commutator is in the middle nucleus of an anti flexible ring and an anti flexible ring without non-zero nilpotent elements is flexible. Using these properties we proved that a semi prime anti flexible ring is a weakly standard ring.

REFERENCES

1. Anderson, C. T. and Outcalt, D. L. "On simple antiflexible rings", Journal of Algebra 10 (1960) 310-320.
2. Kleinfeld, E. "Assosymmetric rings", Proc. Amer. Math. Soc. Vol 8(1957), 983-986.
3. Kosier, F "On a class of nonflexible algebras", Trans. Amer. Math. Soc. 102 (1962), 299-318.

